

Reply to comment by François Bourges *et al.* on “Carbon uptake by karsts in the Houzhai Basin, southwest China”

Junhua Yan,¹ Y. P. Wang,² Guoyi Zhou,¹ Shenggong Li,³ Guirui Yu,³ and Shijie Wang⁴

Received 24 April 2012; revised 21 May 2012; accepted 26 May 2012; published 4 July 2012.

Citation: Yan, J., Y. P. Wang, G. Zhou, S. Li, G. Yu, and S. Wang (2012), Reply to comment by François Bourges *et al.* on “Carbon uptake by karsts in the Houzhai Basin, southwest China”, *J. Geophys. Res.*, 117, G03007, doi:10.1029/2012JG002060.

[1] We thank Bourges *et al.* [2012] for their comment and interest, and we appreciate the opportunity to clarify our paper.

[2] In their commentary, Bourges *et al.* [2012] raise three points: (1) gaseous emissions from the Houzhai Basin are not included in the study of the Houzhai Basin; (2) surface CO₂ emissions as estimated by Bourges *et al.* [2012] are about one order of magnitude greater than the flux estimated by Yan *et al.* [2011]; (3) CO₂ content and aerodynamic regime in the Houzhai Basin can be monitored using a network of flux towers and CO₂ sensors to refine the carbon budget for the basin. In the following we will address each point in turn.

[3] The objectives of the study as outlined in Yan *et al.* [2011] are to understand the temporal variations of carbon uptake from carbonate dissolution and the major driver of that uptake. It was not our objective to construct a full carbon budget for the Houzhai Basin; therefore, gaseous emissions from the land surface were not included in the study (see Figure 1). This addresses point (1).

[4] The carbon budget of most terrestrial ecosystems can be represented by the influx of gross primary production (GPP), efflux of autotrophic and heterotrophic respiration (R_{eco}), and carbon loss from disturbance (L_{dis}), such as fire and wood harvest. The flux of CO₂ dissolved in runoff to rivers or other aquatic systems is usually several orders of magnitude smaller than GPP or R_{eco} and is therefore usually neglected in the carbon budget for most terrestrial ecosystems. However, the runoff CO₂ flux can be a significant component for terrestrial ecosystems in karst landscape because the amount of CO₂ dissolved in karst water can be two orders of magnitude greater than that in a natural water (CO₂ – H₂O) system [Dreybrodt, 1988]. The study [Yan

et al., 2011] estimated that the rate of carbon uptake by chemical weathering in all of karst terrain in China is about 12 Tg C yr⁻¹, which is about 57% of the rate of carbon accumulation by forest biomass in China from 1981 to 1998, suggesting that weathering of carbonate rocks is also an important carbon sink in China. That CO₂ flux via runoff, which depends on soil chemical properties, is sometimes called abiotic flux [Serrano-Ortiz *et al.*, 2010], and it cannot be measured directly by eddy covariance or other aerodynamic techniques. The study by Yan *et al.* [2011] provided one of few estimates for karst systems worldwide.

[5] The CO₂ released from soil respiration in karst terrain can diffuse out to the surface atmosphere via soil pores, macroscopic voids, and fissures, or be chemically taken up during carbonate dissolution. At steady state, the rate of soil respiration is equal to the sum of the rate of CO₂ diffusion to the surface atmosphere and abiotic flux via runoff. Because of the complex air passage in the vadose zone of a karst landscape, the rate of CO₂ diffusion is highly variable (see Figure 1). It is possible that a significant fraction of CO₂ diffusion can occur through macroscopic voids and fissures in a karst landscape, which is what Bourges *et al.* [2012] estimate. Therefore, it is more appropriate to compare the fluxes as estimated by Bourges *et al.* [2012] with the fluxes of ecosystem respiration (R_{eco}) than with the abiotic flux. The flux via diffusion can be much greater than the abiotic uptake via carbonate dissolution, as pointed out by Bourges *et al.* [2012] and others [Kowalczyk and Froelich, 2010; Serrano-Ortiz *et al.*, 2010]. However, these two fluxes can play quite different roles in a regional carbon budget. Abiotic uptake of CO₂ can be transported to rivers in the form of bicarbonate that can be assimilated by some living organisms in aquatic systems [Einsele *et al.*, 2001; Lerman and Mackenzie, 2005; Aufdenkampe *et al.*, 2011]. Once assimilated, much of that carbon will remain in water for decades or even longer, therefore the assimilation of bicarbonate exported from runoff represents a carbon sink, whereas the CO₂ that diffuses out to the surface atmosphere represents a carbon loss from the karst landscape to atmosphere, or a carbon source for the regional carbon budget. This addresses point (2).

[6] Because of many macroscopic voids, numerous fissures, and caves across the Houzhai Basin or similar karst landscape, air passage in the vadose zone is more complicated than in many other landscapes, and surface CO₂ emissions can be extremely variable (see Figure 1). Estimates of CO₂ emission rates from field measurements at the cave opening as recommended by Bourges *et al.* [2012] can be highly uncertain, as

¹Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China.

²CSIRO Marine and Atmospheric Research, Aspendale, Victoria, Australia.

³Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, China.

⁴Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, China.

Corresponding author: J. Yan, Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, 723 Xingke Rd., Tianhe District, Guangzhou 51065, China. (jhyan@scib.ac.cn)




Figure 1. A schematic diagram of the karst landform and CO₂ flow in a karst terrain. The arrows indicate the directions of CO₂ flow. The blue arrows indicate CO₂ emission from macroscopic voids, numerous fissures, and caves, as referred to by Bourges *et al.* [2012]. The red arrows indicate the uptake of CO₂ during rock weathering, and the rate of uptake was estimated by Yan *et al.* [2011] for the Houzhai Basin. The pink arrows indicate the complexity of air passage within the vadose zone in a karst terrain.

the area that contributes to the CO₂ emission at the opening is highly uncertain. We agree that a comprehensive network is required to quantify the carbon budget of the Houzhai Basin using a combination of various techniques. This is the aim of one of our ongoing projects in the Houzhai Basin. This addresses point (3).

[7] **Acknowledgments.** We thank Zaihua Liu, Xiaoyong Bai, and Tao Peng of the Institute of Geochemistry (CAS) for their helpful discussions.

References

- Aufdenkampe, A. K., E. Mayoga, P. A. Raymond, J. M. Melack, S. C. Doney, S. R. Alin, R. E. Aalto, and K. Yoo (2011), Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere, *Ecol. Environ.*, 9(1), 53–60, doi:10.1890/100014.
- Bourges, F., P. Genton, D. Genty, A. Mangin, and D. D'Hulst (2012), Comment on “Carbon uptake by karsts in the Houzhai basin, southwest China” by Junhua Yan *et al.*, *J. Geophys. Res.*, 117, G03006, doi:10.1029/2012JG001937.
- Dreybrodt, W. (1988), *Processes in Karst Systems: Physics, Chemistry, and Geology*, Springer, Berlin, doi:10.1007/978-3-642-83352-6.
- Einsele, G., J. Yan, and M. Hinderer (2001), Atmospheric carbon burial in modern lake basins and its significance for the global carbon budget, *Global Planet. Change*, 30, 167–195, doi:10.1016/S0921-8181(01)00105-9.
- Kowalczyk, A. J., and P. N. Froelich (2010), Cave air ventilation and CO₂ outgassing by radon-222 modeling: How fast do caves breathe?, *Earth Planet. Sci. Lett.*, 289, 209–219, doi:10.1016/j.epsl.2009.11.010.
- Lerman, A., and F. T. Mackenzie (2005), CO₂ air-sea exchange due to calcium carbonate and organic matter storage, and its implications for the global carbon cycle, *Aquat. Geochem.*, 11, 345–390, doi:10.1007/s10498-005-8620-x.
- Serrano-Ortiz, P., M. Roland, S. Sanchez-Moral, I. A. Janssens, F. Domingo, Y. Godderis, and A. S. Kowalski (2010), Hidden, abiotic CO₂ flows and gaseous reservoirs in the terrestrial carbon cycle: Review and perspectives, *Agric. For. Meteorol.*, 150, 321–329, doi:10.1016/j.agrformet.2010.01.002.
- Yan, J., Y. P. Wang, G. Zhou, S. Li, G. Yu, and K. Li (2011), Carbon uptake by karsts in the Houzhai Basin, southwest China, *J. Geophys. Res.*, 116, G04012, doi:10.1029/2011JG001686.