



Evaluation of irrigation effects using diluted salted water based on electrophysiological properties of plants

Qaiser Javed, Yanyou Wu, Ahmad Azeem & Ikram Ullah

To cite this article: Qaiser Javed, Yanyou Wu, Ahmad Azeem & Ikram Ullah (2017) Evaluation of irrigation effects using diluted salted water based on electrophysiological properties of plants, *Journal of Plant Interactions*, 12:1, 219-227, DOI: [10.1080/17429145.2017.1319501](https://doi.org/10.1080/17429145.2017.1319501)

To link to this article: <https://doi.org/10.1080/17429145.2017.1319501>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 05 May 2017.



Submit your article to this journal [↗](#)



Article views: 277



View Crossmark data [↗](#)

RESEARCH ARTICLE



Evaluation of irrigation effects using diluted salted water based on electrophysiological properties of plants

Kaiser Javed^a, Yanyou Wu^b, Ahmad Azeem^a and Ikram Ullah^a

^aKey Laboratory of Modern Agricultural Equipment and Technology, Ministry of Education, Institute of Agricultural Engineering, Jiangsu University, Zhenjiang, People's Republic of China; ^bResearch Center for Environmental Bio-Science and Technology, State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, People's Republic of China

ABSTRACT

Effect of salt stress and subsequent re-watering on growth and electrophysiological characteristics of *Orychophragmus violaceus* and *Brassica napus* was investigated, to construct a model for prediction of an appropriate regime for dilution of saline water. Different concentrations of NaCl, Na₂SO₄ and blend of both salts were applied with Hoagland solution. Growth and electrophysiology traits were highly restricted at high concentration levels. According to the results, the best dilution point was 5–2.5% for NaCl and Na₂SO₄ alternatively, whereas it was 10–0.0% for blend of salts. After re-watering, *O. violaceus* restore better leaf tensity under medium levels, because the effect of NaCl concentration on leaf tensity (X_{NC-LT}) was 1.66%, which was lower in *O. violaceus* than the effect of X_{NC-LT} (2.88%) in *B. napus*, followed by the same trend for all treatments. Therefore, the effect of salinity in *O. violaceus* and *B. napus* may also be reduced effectively by dilution of saline irrigation.

ARTICLE HISTORY

Received 19 January 2017
Accepted 11 April 2017

KEYWORDS

Dilution model; growth; leaf tensity; physiological capacitance; salt stress; re-watering regime

Introduction

In abiotic stresses, drought and salinity are two environmental factors which are responsible for the huge loss in agricultural demand and productivity all over the world (Gul & Ahmad 2004). Water scarcities are expected to lead to loss of crop production globally up to 30% by 2025 compared to current yield. Salinity of agricultural soil is becoming a serious issue due to increasing shortage of fresh water and subsequent necessity to utilize saline water or low-quality water for irrigation (SEMÍZ & Suarez 2015). Therefore, at the current situation of intentions of improving agricultural production under limited water resources, it is required to grow crops that are capable of tolerating drought and water stress environment (Yao et al. 2016). It would be the most economical approach to improve productivity based on limited water resources.

Orychophragmus violaceus and *Brassica napus* from Brassicaceae family are imperative oilseed crops, widely used for human food due to its improved oil content, high nutritious value, enrichment and stability in yield (Purty et al. 2008). Most species of brassica are considered to be moderate salt-tolerant (Masidur Alam et al. 2007). However, their growth and yield were markedly reduced due to stresses (Su et al. 2013) and also the production of edible oil crops was affected at a distressing rate due to rising issues of water and soil salinization (García-Sánchez et al. 2002). In order to provide attention to these aspects of these crops, studies were necessary according to variation in the morphological characteristics. Later on the experiment was conducted to emphasize on plants physiognomies under water and salt stress (Zhang et al. 2012; Benincasa et al. 2013; de Oliveira et al. 2013).

In the field, plants may have experienced several diverse abiotics, either contemporarily at different times throughout

the growing season (Tester & Bacic 2005). Exposure of plants to a traumatic environment during development periods causes many physiological changes (Jamil et al. 2012). The most common adverse effect of salt stresses on the physiology is the reduction in plant growth and yield (Ashraf & McNeilly 2004; Abdul Qados 2011). Growth is the most important process inhibited, partially or completely by stresses. However, irrigation water quality may have a profound impact on plant growth including the series of processes. The first response of plants towards salinity stress is a reduction of leaf growth. If salt stress ultimately rises to toxic levels in the leaves, it initiates premature senescence and reduces leaf area to a level that cannot withstand growth characteristics (Munns 2002). Toxic effect of salt stress repressed leaf expansion (Roy et al. 2014) and reduces the production of new leaves (Mansour & Salama 2004). Salt tolerance in plants may be explained by functional and structural adaptations, such as growth regulation, water potential and osmotic adjustment. Osmotic adjustment is an adaptation mechanism for the plant to maintain their water stability (Zollinger et al. 2007).

Water stability and deficiency in plants can also be determined using their electrophysiological properties. In which, the water potential of a plant was reflected by water content, leaf air temperature and capacitance (Koide et al. 1989). Water potential describes the energy level of water movement (Vogt & Losch 1999). A significant relationship is observed between water content, water potential and physiological capacitance. Therefore, water status in leaves is found from the variations of physiological resistance and capacitance. The status of water in plants is also characterized by the tensity of the plant cell, which reflects the salts resistance of plants (Zhang et al. 2015). To overwhelm this situation, salt stress, subsequent re-watering or dilution of saline irrigation

could be helpful for plants to recover their growth development through rebuilding of leaf tensity, maintain water potential and water content.

Prediction of re-watering or the dilution point of saline water is a new index, which could be helpful for the regulation of saline water, in order to sustain agricultural productivity and economical irrigation. An appropriate dilution of salt water on an appropriate point will save the water resource. *O. violaceus* and *B. napus* are hypothesized for better restorability after re-watering or dilution of salted water. Therefore, the intention of this present study was to observe the electrophysiological and growth responses of *O. violaceus* and *B. napus* under different salt stresses and subsequent re-watering conditions. Furthermore, to optimize plant growth and yield, a model needs to be constructed for predicting an appropriate regime for dilution of saline water on the basis of electrophysiological and growth responses.

Materials and methods

Experimental conditions

The experiment was carried out at the Institute of Agricultural Engineering, Jiangsu University, Zhenjiang, Jiangsu, China (32.20°N, 119.45°E). Intact seeds of *O. violaceus* and *B. napus*, which were identical in size and color, homogeneous and free from wrinkles, were chosen for this experiment. Seeds were cultivated in 20-cell trays, containing distilled water and washed vermiculite in equal amount. The seeds were left to grow inside the growth chamber under day/night temperature cycle of 25°C/20°C, and 60% of relative humidity.

Salt treatments

Plants were irrigated daily with Hoagland solution (Hoagland & Arnon 1950). After 20 days, plants were transferred to the greenhouse under natural lighting, (25/18) ± 2°C (day/night) and 70% relative humidity. Homogenous seedlings, showing healthy growth were selected for further treatments. They were exposed to salt stress, induced by NaCl, Na₂SO₄ and combination of both salts at different four levels, in which one is the control level. The concentration of NaCl (NC1:2.5, NC2:5.0, NC3:10.0 and 0 as control) g L⁻¹; Na₂SO₄ (NS1:2.5, NS2:5.0, NS3:10.0 and 0 as control) g L⁻¹ and in blend of salts (BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control) were selected for treatments and applied to plants with Hoagland solution. The control plants only received a nutrient solution.

Re-watering orders

Re-watering was continued on day 14 from the onset of salt stress treatment. The order for re-watering was that, plants that were affected by high stress level (10.0 g L⁻¹ in both NaCl and Na₂SO₄) were irrigated with medium stress level NC3², NS3² (5.0 g L⁻¹ in both NaCl and Na₂SO₄, respectively), afterwards medium stress level (5.0 g L⁻¹ in both NaCl and Na₂SO₄) irrigated with low stress level NC2¹, NS2¹ (2.5 g L⁻¹ in both NaCl and Na₂SO₄, respectively) and then low stress level (2.5 g L⁻¹ in both NaCl and Na₂SO₄, respectively) irrigated with control NC1⁰, NS1⁰ (0 g L⁻¹ in both

NaCl and Na₂SO₄, respectively). In mixed treatments, all levels were re-watered with control (BS1⁰, BS2⁰, BS3⁰). This experiment was designed in the randomized block design and a total of three replicates was chosen for each physiological measurements.

Determination of growth parameters

Growth measurements were considered after treatments and after re-watering three times per each week in both cases, respectively. The five replicates were chosen for each treatment and also used to analyze the mean of each measurement. The measurements taken for growth analysis were: plant height (PH); stem diameter (SD) and leaf area (LA). The leaf area was measured by a leaf area meter (a handheld laser leaf area meter, CI, 203).

Determination of leaf water potential and leaf water content

Leaves in salt stress phase and in subsequent re-watering phase were used for the determination of water potential (Ψ) and water content (WC). Leaves' water potential were measured at 9:00–11:00 am with a dew point microvolt meter in a C-52-SF universal sample room (Psypro, Wescor, USA), every 3 days in both salt stress and re-watering phases, respectively. Five plants from each treatment group were selected for the measurement. The water content in leaves of *O. violaceus* and *B. napus* was measured through the relationship between the water potential (Ψ) of a plant tissue and cell SAP solute concentration (Q), and is

$$\Psi = iQRT, \quad (1)$$

where Ψ is the water potential (MPa) of a plant tissue; *I* the dissociation coefficient (*i* = 1); *Q* the cell SAP solute concentration (mol L⁻¹); *R* the gas constant (*R* = 0.0083 L MPa mol⁻¹ K⁻¹) and *T* the thermodynamic temperature (273 + *t* °C) K.

The relationship between proportion of solute quality in the total quality of leaf (*P*, %) and cell SAP solute concentration (*Q*) is

$$P = \frac{MQ}{1000} \%, \quad (2)$$

where *M* is the relative molecular mass of cell SAP solute, sugar C₁₂H₂₂O₁₁, *M* was 342 g mol⁻¹.

According to Equations (1) and (2), *P* could be rewritten as follows:

$$P = \left(-\frac{\Psi M}{100iRT} \right) \%. \quad (3)$$

The proportion of water content in the total quality of leaf was 1 - *P*; it was expressed as WC (%)

$$WC = \left(1 + \frac{\Psi M}{100iRT} \right) \%. \quad (4)$$

Determination of leaf tensity

The leaf was clipped between two electrodes of the parallel plate capacitor, which formed a parallel plate-capacitor sensor. The water potential of the leaves changed with variations

in cytosol solute concentration in the leaves. This behavior caused a change in the dielectric constant of the cytosol solute in the leaf tissue that was between the two electrodes. As a result, the physiological capacitance (C_p) of the plant was affected.

The relationship between the proportion of solute quality in the total quality of the leaf (P) and the concentration (Q) was $Q = 1000 P/M$ where M is the relative molecular mass of the cytosol solute. Comparing the relationship between P , Q , and C_p , the values of leaf tensity (T_L) is rewritten according to (Zhang et al. 2015) as shown below:

$$\frac{d}{A} = \frac{\epsilon_o \cdot \left[81 - \frac{(81 - a)MW}{1000iRT} \right]}{d} \quad (5)$$

and

$$y = \frac{d}{A} = \frac{\epsilon_o}{C_p} \left[81 + \frac{(81 - a)MW}{1000iRT} \right]. \quad (6)$$

For a given material, the relative dielectric constant ($a = 3.3$) and the molecular mass ($M = 342$) of the cytosol solute were all assigned values. The leaf tensity of the plants [$T_L = 1/y$ ($i = 0, 1, 2, 3$)] was determined to indicate the water status in the leaf. In this study, sugar $C_{12}H_{22}O_{11}$ was identified as the solute in the cytosol; therefore, T is (K, $T = 273 + t$ °C), R is the gas constant ($R = 0.0083$ L MPa mol⁻¹ K⁻¹) and i is the dissociation coefficient equal to 1.

Model construction for prediction of re-watering levels

The exponential decay equation is used to construct the model between salt stress levels and C_p , also to construct the model between salt stress levels and T_L , these are shown in Equations (5) and (6). Here, a_1, a_2, b_1, b_2 are constants and X_{EC} is the salt concentrations in the form of a percentage (%).

$$C_p = a_1 e^{-b_1 X_{EC}}, \quad (7)$$

$$T_L = a_2 e^{-b_2 X_{EC}}. \quad (8)$$

Predictions of re-watering levels were calculated using Equations (7) and (8). Substituting the values of C_p and T_L which were obtained through after re-watering into Equations (7) and (8) gave the point of predicted re-watering levels, equivalent to salt stress levels.

Statistical analysis

All measurements were subjected to analysis of variance (ANOVA) to discriminate significant differences (defined as $p \leq 0.05$) between group means. Data are shown as the mean \pm standard error (SE) ($n = 5$). These mean data were analyzed statistically using a factorial design through SPSS software (version 13.0, SPSS Inc.), and mean results were compared through LSD at 5% significance level ($p < 0.05$).

Results

Effect of salt stress and re-watering on plant growth

The plant height (PH), stem diameter (SD) and leaf area (LA) of both species showed a noticeable decrease on being exposed to different concentrations of both salts (Tables 1–3). Of the

different concentration of both salts, the lowest concentration (NC1 and NS1) showed a minimum reduction compared with the control. But, the highest concentration of both single salts and blend of both salts at NC3, NS3, BS1, and BS2 made severe reduction in PH, SD and LA, respectively.

During treatment periods of *O. violaceus* at the BS1-concentration level, it triggered a maximum decrease in PH (5.70 cm), SD (0.04 cm) and LA (11.35 cm²), compared with *B. napus*. However, the high concentration of blend of salts caused a more prominent decrease in the growth parameters of *O. violaceus* than *B. napus* as compared to single salts. Moreover, single Na₂SO₄ was found to be toxic to *O. violaceus* and *B. napus* at NS3 than NC3, high concentration level of NaCl, respectively (Tables 1–3).

A general trend appeared in the increments of plant growth in both species during re-watering periods. A statistical analysis specified that the increments which were observed during re-watering was significant, except for the concentration at high levels (NC3[^]2, NS3[^]2, BS1[^]0 and BS2[^]0). The increments in PH, SD and LA were obtained more under the medium concentration of single and blend of salts (NC2[^]1, NS2[^]1 and BS3[^]0) followed by the low concentration of salt (NC1[^]0 and NS1[^]0) as compared to control, respectively. But relatively, the increments of PH (58.27%, 56.20% and 55.95%), SD (62.16%, 60% and 60%) and LA (46.53%, 44.91% and 43.78%) were observed more in *O. violaceus* under NC2[^]1, NS2[^]1 and BS3[^]0, medium concentration of single and blend of salts, compared with the responses of *B. napus* under medium concentration of single and blend of salts, respectively. However, at a high concentration of single and blend of salts (NC3[^]2, NS3[^]2, BS1[^]0 and BS2[^]0), the adverse effect of salt stress on PH, SD and LA, was exhibited. Although, the addition of salts at BS3[^]0 revealed the same effect on PH, SD and LA as compared to PH, SD and LA under NC2[^]1 and NS2[^]1 concentration levels, respectively (Tables 1–3). In general, statistical analysis demonstrated that after re-watering *O. violaceus* showed better growth development than *B. napus*, respectively.

Water content and water potential

Results shown in Tables 4 and 5 indicated that an increase in the concentration of salts decreased the water potential (Ψ) and water content (WC) of leaves in both species during treatments periods. The maximum loss in WC was observed in both species at high concentration levels of both single and blend of salts (NC3, NS3, BS1 and BS2) as compared to control. While the minimum decrease in both WC and Ψ was noted at under low concentration levels (NC1 and NS1) as compared with the control in both *O. violaceus* and *B. napus*, respectively. However, the maximum decrease in Ψ was noted at NC3, NS3, BS1 and BS2, high concentration levels, respectively. While in medium (NC2, NS2 and BS3) concentration of NaCl, Na₂SO₄ and blend of salts, the decrease in WC and in Ψ was found to be almost the same in both *O. violaceus* and *B. napus*, respectively. As a comparison between single salts and a mixture of salt, it was observed that WC and Ψ of *B. napus* were significantly less affected by NaCl concentrations than Na₂SO₄ and blend of salt concentrations. However, the high concentration blend of salts affects *O. violaceus* compared with single salts and caused a more prominent decrease in the WC and Ψ compared with *B. napus*.

Table 1. Effect of salt stress and re-watering on plant height of *O. violaceus* and *B. napus*.

Species Treatments/re- watering (g L ⁻¹)	<i>O. violaceus</i>			<i>B. napus</i>		
	Plant H. under salt stress (cm)	Plant H. under re- watering (cm)	Increment in Plant H. after re-watering (%)	Plant H. under salt stress (cm)	Plant H. under re- watering (cm)	Increment in Plant H. after re-watering (%)
Control	35.30 ± 0.70a	50.40 ± 0.90a	42.78	40.10 ± 0.80a	55.25 ± 1.25a	37.78
NC1/NC1^0	30.45 ± 0.15b	45.05 ± 0.05a	47.95	31.25 ± 0.75b	45.15 ± 1.15b	44.48
NC2/NC2^1	19.05 ± 0.55d	30.15 ± 0.15c	58.27	24.50 ± 0.50c	36.75 ± 0.25c	50.00
NC3/NC3^2	9.50 ± 0.50e	11.40 ± 0.40e	20.00	17.50 ± 0.50d	20.15 ± 0.65d	15.14
NS1/NS1^0	28.45 ± 0.25c	41.50 ± 0.90b	45.87	29.50 ± 0.50b	44.00 ± 2.00b	49.15
NS2/NS2^1	18.15 ± 0.15d	28.35 ± 0.25d	56.20	23.50 ± 0.50c	35.00 ± 0.00c	48.94
NS3/NS3^2	07.85 ± 0.45f	09.25 ± 0.25ef	17.83	14.85 ± 0.85e	17.15 ± 0.15e	15.49
BS1/BS1^0	05.70 ± 0.50g	06.20 ± 1.20f	08.77	10.95 ± 1.00f	11.50 ± 0.50f	05.02
BS2/BS2^0	06.95 ± 0.05fg	08.00 ± 1.00f	15.11	13.15 ± 0.15e	14.70 ± 0.70e	11.79
BS3/BS3^0	18.50 ± 0.30d	28.85 ± 0.55d	55.95	23.05 ± 0.55c	34.85 ± 1.25c	51.19

Notes: The means ± SE (n = 5) followed by different letter in the columns indicated the plant height of both species under salt stress and increment in plant height after re-watering with significant difference at $p \leq .05$, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0: 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 10.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control.

In the re-watering phase, the outcome of the results showed that WC and Ψ of both *O. violaceus* and *B. napus* were recovered (Tables 4 and 5). The increment in WC and Ψ increased under low (NC1^0 and NS1^0) to medium concentration levels (NC2^1, NS2^1 and BS3^0). But relatively, the increments of WC (10.41%, 12.50% and 12.40%) were observed more in *O. violaceus* under NC2^1, NS2^1 and BS3^0, medium concentration of single and blend of salts, compared with the results of *B. napus* (4.79%, 7.38% and 9.22%) under medium concentration of single and blend of salts, respectively.

But the degree of salts still showed adverse effects at high concentration levels (NC3^2, NS3^2, BS1^0 and BS2^0) with an increment of WC and Ψ even during the re-watering phase. Consequently, the blend of salts at BS3^0 revealed the same effect on WC and Ψ as compared with WC and Ψ under NC2^1 and NS2^1 concentration levels, respectively. Overall, the statistical analysis revealed that *O. violaceus* showed better restorability of WC and Ψ after re-watering than *B. napus*, respectively.

Physiological capacitance and leaf tensity

From the study, it was found that the physiological capacitance (C_p) and leaf tensity (T_L) significantly decreased with increasing salt stress in comparison with the control (Tables

6 and 7) in both species. *O. violaceus* showed better tolerance than *B. napus* from low (NC1 and NS1) to medium (NC2, NS2 and BS3) concentration of NaCl, Na₂SO₄ and blend of salts. Maximum reduction of C_p and T_L was noted in both *O. violaceus* and *B. napus* under high concentrations at NC3, NS3, BS1 and BS2 levels of NaCl, Na₂SO₄ and blend of salts, respectively, as compared with control. While the minimum reduction of C_p and T_L was recorded in both *O. violaceus* and *B. napus* under low (NC1 and NS1) concentrations of NaCl and Na₂SO₄ as compared with C_p and T_L under control condition. *O. violaceus* exhibited the highest reduction in C_p and T_L under BS1 concentration of a blend of salts as compared with *B. napus*. However, in medium (NC2, NS2 and BS3) concentration of NaCl, Na₂SO₄ and mixture of salts, the reduction in C_p and T_L was found to be almost the same.

The physiological capacitance (C_p) and leaf tensity (T_L) significantly increased during salt stress followed by re-watering. It was observed that *O. violaceus* exhibited better results compared with *B. napus* from the stress phase to the re-watering phase. During re-watering, C_p and T_L increased under treatments with low concentration levels of NC1^0 and NS1^0, respectively. However, salt stress in both species at high concentration affected C_p and T_L adversely following the order NC3^2, NS3^2, BS1^0 and BS2^0, respectively. While in the re-watering phase, the increment in T_L under low salt stress treatments (NC1^0 and NS1^0) was found

Table 2. Effect of salt stress and re-watering on stem diameter of *O. violaceus* and *B. napus*.

Species Treatments/re- watering (g L ⁻¹)	<i>O. violaceus</i>			<i>B. napus</i>		
	Stem D. under salt stress (cm)	Stem D. under re- watering (cm)	Increment in Stem D. after re-watering (%)	Stem D. under salt stress (cm)	Stem D. under re- watering (cm)	Increment in Stem D. after re-watering (%)
Control	0.33 ± 0.01a	0.49 ± 0.01a	48.48	0.56 ± 0.01a	0.70 ± 0.00a	25.00
NC1/NC1^0	0.28 ± 0.01b	0.43 ± 0.00b	51.79	0.50 ± 0.01b	0.65 ± 0.01b	30.00
NC2/NC2^1	0.19 ± 0.01c	0.30 ± 0.00c	62.16	0.44 ± 0.01c	0.54 ± 0.01c	22.99
NC3/NC3^2	0.09 ± 0.01d	0.10 ± 0.00d	17.65	0.31 ± 0.01d	0.34 ± 0.02d	09.84
NS1/NS1^0	0.26 ± 0.01b	0.40 ± 0.00b	53.85	0.51 ± 0.01b	0.65 ± 0.01b	27.45
NS2/NS2^1	0.18 ± 0.00c	0.28 ± 0.01c	60.00	0.43 ± 0.01c	0.52 ± 0.01c	21.18
NS3/NS3^2	0.08 ± 0.01de	0.09 ± 0.02de	13.33	0.29 ± 0.00de	0.32 ± 0.01de	08.62
BS1/BS1^0	0.04 ± 0.01e	0.04 ± 0.01e	08.57	0.29 ± 0.00e	0.30 ± 0.01e	03.51
BS2/BS2^0	0.06 ± 0.01e	0.06 ± 0.01e	09.09	0.29 ± 0.00de	0.31 ± 0.01e	06.90
BS3/BS3^0	0.18 ± 0.00c	0.28 ± 0.01c	60.00	0.54 ± 0.00c	0.54 ± 0.01c	25.58

Notes: The means ± SE (n = 5) followed by different letter in the columns indicated the stem diameter of both species under salt stress and increment in stem diameter after re-watering with significant difference at $p \leq 0.05$, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0: 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 10.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control.

Table 3. Effect of salt stress and re-watering on leaf area of *O. violaceus* and *B. napus*.

Species Treatments/re- watering (g L ⁻¹)	<i>O. violaceus</i>			<i>B. napus</i>		
	Leaf A. under salt stress (cm)	Leaf A. under re- watering (cm)	Increment in Leaf A. after re-watering (%)	Leaf A. under salt stress (cm)	Leaf A. under re- watering (cm)	Increment in Leaf A. after re-watering (%)
Control	48.75 ± 1.25a	55.50 ± 2.00a	21.31	55.57 ± 1.18a	70.60 ± 0.96a	27.05
NC1/NC1^0	40.25 ± 0.25b	54.00 ± 1.00a	34.16	46.51 ± 0.97b	60.32 ± 0.04b	29.68
NC2/NC2^1	28.80 ± 0.30c	42.20 ± 0.60b	46.53	40.63 ± 0.34d	55.10 ± 0.54c	35.63
NC3/NC3^2	17.90 ± 0.90d	21.00 ± 1.00c	17.32	30.50 ± 0.50e	35.25 ± 0.75d	15.57
NS1/NS1^0	39.10 ± 1.10b	53.05 ± 0.75a	35.68	43.50 ± 0.50c	59.39 ± 1.03b	36.53
NS2/NS2^1	28.50 ± 0.50c	41.30 ± 1.30b	44.91	39.05 ± 1.01d	52.84 ± 1.84c	35.31
NS3/NS3^2	16.30 ± 0.40d	18.70 ± 0.10c	14.72	28.69 ± 1.57ef	31.78 ± 0.77e	10.75
BS1/BS1^0	11.35 ± 1.05e	12.00 ± 1.00d	05.73	26.00 ± 0.00f	27.25 ± 0.25f	04.81
BS2/BS2^0	12.55 ± 0.55e	13.55 ± 0.45d	07.97	27.69 ± 0.57f	29.44 ± 0.44ef	06.32
BS3/BS3^0	28.55 ± 0.05c	41.05 ± 0.35b	43.78	38.23 ± 0.05d	52.49 ± 0.49c	37.32

Notes: The means ± SE ($n = 5$) followed by different letter in the columns indicated the leaf area of both species under salt stress and increment in leaf area after re-watering with significant difference at $p \leq .05$, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0: 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 0.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control.

to be the highest at 0.106 and 0.102 cm in *O. violaceus*, 0.113 and 0.113 cm in *B. napus*, respectively. Relatively, more increments of C_p and T_L were found in *O. violaceus* under NC2^1, NS2^1 and BS3^0, medium concentration levels, respectively. However, *O. violaceus* showed better recovery of C_p and T_L from low to medium concentration levels than *B. napus* (Tables 6 and 7).

Model construction relationship between leaf area and physiological capacitance

By using data of C_p and T_L under salt stress levels, a model using exponential decay equations can be given as shown in Table 8.

It is noted that the relationship between salt stress concentrations (X_C) and C_p or leaf tensity and between X_C and T_L can characterize well the relationship between leaf water status and salt stress concentrations. The coefficient of determination (R^2) showed that the relationship between C_p and X_C in *O. violaceus* fits better with the exponential decay equation as compared with the relationship between T_L and X_C . While in *B. napus*, R^2 showed that the relationship between C_p and X_C is the same with the exponential decay equation as compared with the relationship between T_L and X_C .

Re-watering or dilute irrigation points were almost the same by using the C_p and T_L equation for both species. The values of dilute irrigation of both cultivars are shown in Table 9.

But after dilution, *B. napus* showed sensitive behavior as compared to *O. violaceus*. According to C_p and T_L model equations, it is cleared that *B. napus* and *O. violaceus* have to re-water or dilution of salted water when X_C was reached to 5%. When the plant was exposed to 10% of X_C in the case of single NaCl (X_{NC}) and Na₂SO₄ (X_{NS}) and 12.5% in the case of the blend of salts (X_{MNC} and X_{MNS}) more reduction in plant growth was found (Tables 1–3). Consequently, the effect of dilution on C_p and T_L under 5–2.5% concentration was 1.66% (X_{NC-LT}), 1.74% (X_{NC-CP}), 1.45% (X_{NS-LT}), 1.28% (X_{NS-CP}), 1.60% (X_{MNC-LT}), 1.94% (X_{MNC-CP}), 1.45% (X_{MNS-LT}) and 1.77% (X_{MNC-CP}), respectively, in *O. violaceus* and 2.88% (X_{NC-LT}), 2.98% (X_{NC-CP}), 2.92% (X_{NS-LT}), 3.01% (X_{NS-CP}), 3.15% (X_{MNC-LT}), 2.95% (X_{MNC-CP}), 2.97% (X_{MNS-LT}) and 2.81% (X_{MNC-CP}), respectively, in *B. napus*. Therefore, recovery was more in *O. violaceus*

under 5–2.5%, because of dilution effects, observed on plants was 1.66% of X_{NC-LT} but in *B. napus* the effect was observed 2.88% of X_{NC-LT} followed by the same trend for all the concentrations. Hence, this regime is considered the best for dilution of salted water for both the species. Therefore, the dilution point is the best condition to start re-watering or dilution of saline water for plant growth development because beyond that point plants will be suffer serious stress.

Discussion

Effect of salt stress on electrophysiological properties

O. violaceus and *B. napus* have varied inconsistency in their response to salt stress. The responses of the plants to salt stresses in terms of growth development are definitive expressions of many physiological processes. A surprising concentration of salts affected the capability of roots to extract water, resulting in the limitation of many physiological processes such as hindrance in nutrients uptake (Munns & Tester 2008) and reduction in the plant growth and development (Zapryanova & Atanassova 2009; Wang et al. 2011). Stress caused excessive reduction in PH, SD and LA at high stress levels (NC3, NS3, BS1 and BS2), due to leaf tissue dehydration and limiting photosynthesis activity through stomatal closure (Tables 1–3). At that point, there was a negative relationship between plant growth and salts' concentration, in which the salts' accumulation in leaves directly affect Ψ (Niu et al. 2008; Fu et al. 2010) by decreasing the water content of the leaves. However, the lowest Ψ values were recorded in plants when subjected to water stress (Álvarez et al. 2012).

Salt stress at high concentration levels in both species exerts its effect directly on C_p by disturbing the water status of leaves, which was in response to the effect on leaf cell extension. The C_p of *O. violaceus* was decreased and affected more as compared to *B. napus* at high concentration levels of single and blend of salts and Ψ also decreased, which indicates that water absorption ratios were disturbed due to leaf cell shrinkage. The decrease in leaf area was a response to the compact expansion of the leaf cell. However, minimum reduction in C_p of *O. violaceus* and *B. napus* from low (NC2 and NS2) to medium concentration levels (NC2, NS and BS3) indicates that the water-holding capability of the leaf cell might have played a vital role to maintain leaf WC.

Table 4. Effect of salt stress and salt stress subsequently re-watering on water potential of *O. violaceus* and *B. napus*.

Treatments/re-watering (g L ⁻¹)	$\Psi_{O. violaceus}$			$\Psi_{B. napus}$		
	Salt -stress (MPa)	Re-watering (MPa)	Increment (MPa)	Salt -stress (MPa)	Re-watering (MPa)	Increment (MPa)
Control	-0.95 ± 0.03a	-0.91 ± 0.03a	0.05	-1.19 ± 0.06a	-0.90 ± 0.09a	0.29
NC1/NC1^0	-1.70 ± 0.08b	-1.20 ± 0.02b	0.50	-1.23 ± 0.13ab	-1.01 ± 0.13b	0.22
NC2/NC2^1	-2.19 ± 0.09c	-1.51 ± 0.14c	0.68	-1.82 ± 0.06ab	-1.56 ± 0.01b	0.26
NC3/NC3^2	-3.24 ± 0.05d	-3.06 ± 0.05e	0.18	-3.05 ± 0.25c	-2.94 ± 0.09c	0.11
NS1/NS1^0	-1.74 ± 0.05b	-1.26 ± 0.03b	0.48	-1.46 ± 0.23ab	-1.07 ± 0.10b	0.39
NS2/NS2^1	-2.34 ± 0.07c	-1.73 ± 0.04d	0.61	-2.14 ± 0.10b	-1.77 ± 0.09b	0.38
NS3/NS3^2	-3.35 ± 0.04d	-3.19 ± 0.11e	0.16	-3.79 ± 0.97cd	-3.66 ± 0.97c	0.14
BS1/BS1^0	-4.58 ± 0.06e	-4.45 ± 0.10g	0.13	-4.58 ± 0.58d	-4.53 ± 0.57d	0.05
BS2/BS2^0	-4.20 ± 0.04f	-4.03 ± 0.02f	0.17	-3.77 ± 0.15c	-3.71 ± 0.16c	0.06
BS3/BS3^0	-2.36 ± 0.08c	-1.75 ± 0.04d	0.61	-1.75 ± 0.14ab	-1.25 ± 0.23b	0.51

Notes: The means ± SE ($n = 5$) followed by different letter in the columns indicated the water potential of both species under salt stress and increment in water potential after re-watering with significant difference at $p \leq .05$, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0: 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 10.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control.

However, the decrease and increase in the physiological capacitance of the leaves indicate the concentration of solute in the cytosol. The decrease in the concentration of solute in the cytosol describes the threshold cell's value for dysfunction. Meanwhile the increase and decrease in Ψ indicate the plant's physiological method for survival when the loss of water is continued (Zhang et al. 2015). Meinzer et al. (2003) use the values of C_p to measure water transport properties in trees of sapwood. As a whole, there is need to determine Ψ and C_p together at the same time because both reflect the leaf water status of plants. The synergetic effect of Ψ and C_p allows the plants to survive under water stress environment.

Leaves' cell tension has the ability to sustain water in order to maintain development of plants. Leaf cell tension is used to describe the plant's water status. The variations in tension reflect the amount of water lost and the ability of the leaf cell to hold water (Irigoyen et al. 1992). The capability of holding water is directly linked with the drought tolerance of plants under stress conditions. The results in Table 7 show that the minimum reduction found from low-medium stress levels in T_L of *B. napus* indicates the minimum change in T_L and the ability of *B. napus* to hold water. While the changes in T_L of *O. violaceus* were also found to be minimum under low-medium stress levels, but water-holding capability of *O. violaceus* was not better as found in *B. napus*. However at high stress levels, T_L of *O. violaceus* and *B. napus* reduced

sharply and their ability to hold water was not good. Thus, this condition exposed the threshold flexibility and tolerance of *O. violaceus* and *B. napus* under low-moderate levels because of the mechanisms involved in osmotic tolerance related to water availability to sustain leaf cell's energy demands for growth.

Re-watering or dilution point

Predicted re-watering salt stress levels based on C_p and T_L showed that *O. violaceus* and *B. napus* have to re-water when the plant goes under medium concentration levels (Table 8). In the absence of salt stress, predicted values for both species were near to zero, indicating that the present finding is credible. For *O. violaceus*, in addition to the control (no salt stress), the medium stress levels of NaCl and Na₂SO₄ from 5.0% rehydration to 2.5% treatment and from 10% rehydration to 0.0% treatment in blend of salts indicate that after dilution, adaptability of plants is greatly enhanced. Later, adaptability of *B. napus* is greatly enhanced, but not as observed in the case of *O. violaceus*. The results of growth parameters of both species were also found to be better in predicting these re-watering levels (Tables 1–3), indicating that these predicting re-watering levels techniques are very credible. Best predicting re-watering levels and dilution points of both species were also found in the same regime that also strengthens our findings. Because beyond that point up

Table 5. Effect of salt stress and salt stress subsequently re-watering on water content of *O. violaceus* and *B. napus*.

Treatments/re-watering (g L ⁻¹)	WC _{O. violaceus}			WC _{B. napus}		
	Salt -stress (%)	Re-watering (%)	Increment (%)	Salt -stress (%)	Re-watering (%)	Increment (%)
Control	86.85 ± 0.47a	87.50 ± 0.40a	0.74	83.58 ± 0.86a	86.85 ± 0.49a	3.91
NC1/NC1^0	76.50 ± 1.15b	83.40 ± 0.30b	9.01	83.03 ± 1.81a	86.25 ± 1.66a	3.87
NC2/NC2^1	69.75 ± 1.25c	77.01 ± 0.41c	10.41	74.85 ± 0.88a	78.44 ± 0.20a	4.79
NC3/NC3^2	55.26 ± 0.69d	57.79 ± 0.65d	4.58	57.88 ± 2.49b	59.11 ± 1.09b	2.12
NS1/NS1^0	76.00 ± 0.65b	82.62 ± 0.44b	8.72	79.91 ± 2.16a	85.29 ± 1.45a	6.73
NS2/NS2^1	67.68 ± 0.94c	76.14 ± 0.52c	12.50	70.43 ± 1.34ab	75.63 ± 1.22a	7.38
NS3/NS3^2	53.84 ± 0.56d	56.04 ± 1.54d	4.10	47.67 ± 3.41b	49.56 ± 3.36b	3.96
BS1/BS1^0	32.50 ± 0.87f	34.02 ± 1.44f	4.67	32.09 ± 3.05c	32.73 ± 2.83c	2.02
BS2/BS2^0	41.38 ± 0.52e	43.67 ± 0.28e	5.55	47.95 ± 2.06b	48.83 ± 2.24b	1.82
BS3/BS3^0	67.45 ± 1.07c	75.81 ± 0.59c	12.40	75.82 ± 1.95a	82.80 ± 1.18a	9.22

Notes: The means ± SE ($n = 5$) followed by different letter in the columns indicated the water content of both species under salt stress and increment in water content after re-watering with significant difference at $p \leq .05$, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0 : 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 10.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control.

Table 6. Effect of salt stress and salt stress subsequently re-watering on physiological capacitance of *O. violaceus* and *B. napus*.

Treatments/ Re-watering (g L ⁻¹)	<i>C_{PO}. violaceus</i>		<i>C_{PB}. napus</i>	
	Salt stress (pF)	Re-watering (pF)	Salt stress (pF)	Re-watering (pF)
Control	75.60 ± 1.68a	82.77 ± 1.97a	94.03 ± 0.88a	97.87 ± 0.38a
NC1/NC1^0	52.07 ± 1.42b	63.60 ± 1.89b	63.57 ± 2.88b	70.33 ± 0.43b
NC2/NC2^1	38.40 ± 0.95c	57.93 ± 1.55c	40.73 ± 1.27d	54.83 ± 2.52d
NC3/NC3^2	09.19 ± 0.07e	15.60 ± 0.10e	6.84 ± 0.14f	12.90 ± 0.21f
NS1/NS1^0	48.67 ± 0.90b	61.00 ± 1.63bc	58.60 ± 1.15c	67.10 ± 1.77c
NS2/NS2^1	36.50 ± 0.87c	56.03 ± 0.84cd	36.80 ± 1.66e	51.00 ± 1.31e
NS3/NS3^2	09.01 ± 0.02e	15.50 ± 0.05e	05.45 ± 0.13f	10.60 ± 0.17f
BS1/BS1^0	04.91 ± 0.03e	08.60 ± 0.62e	03.20 ± 0.31f	07.69 ± 0.44f
BS2/BS2^0	06.29 ± 0.03e	10.50 ± 0.81e	05.06 ± 0.09f	09.97 ± 0.07f
BS3/BS3^0	24.84 ± 2.56d	52.67 ± 2.12d	38.67 ± 0.11de	53.13 ± 1.19de

Note: The means ± SE (n = 5) followed by different letter in the columns indicated the physiological capacitance of both species under salt stress followed by re-watering with significant difference at P ≤ .05, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0: 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 10.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control.

to high stress levels of single NaCl and Na₂SO₄ and blend of salts, growth was inhibited. When the plant was exposed to high salinity levels, 50% reduction is found in plant growth (Siringam et al. 2011, Aref 2013). Our results for *O. violaceus* and *B. napus* also agreed with their finding, because both species showed sudden reduction in plant growth at high stress levels (NC3, NS3, BS1 and BS2), and there was no recovery found when X_{NC} and X_{NS} were diluted to 10–5% and X_{MNC} and X_{MNS} were diluted from 12.5% to 0.0%, respectively (Table 9). Brodribb and Holbrook (2005) explained that at 50% collapse of cells, the plant could not recover well during re-watering. Therefore, because of this reason 30% yield loss can be considered acceptable for re-watering in this study. According to 70% increment in plant growth during salt stress, medium concentration levels (NC2, NS2 and BS3) of NaCl, Na₂SO₄ and blend of salts is considered the best for the production of *B. napus* under salt-stressed conditions. But in the case of *O. violaceus*, it exposed the sensitive behavior but after re-watering or dilution, it showed better restoration. It might be that after re-watering of *B. napus*, C_P and T_L under medium to high concentration levels did not increase much and solute

concentration in the cytosol could have reached below the threshold value due to continuous stress which destroyed the leaf cells (Zhang et al. 2015).

From the results (Table 9), it is clear that the dilution point (5–2.5%) under NaCl concentrations was the same for both species which were 1.66% (X_{NC-LT}) and 2.88% (X_{NC-LT}) for *O. Violaceus* and *B. napus*, respectively. Because at 10% salt concentration, salt stress was too severe to be re-watered, and the irrigation effects using diluted salted water (2.5%) at 5% salt stress level in *O. violaceus* and *B. napus* were equivalent to the effects using 1.66% salted water and 2.88% salted water, which were equal to the salt stress level between 0% and 2.5%. Accordingly, when the concentration of salted water was 2.5% in both species, it could be used for irrigation directly, but when the concentration of salted water was 5% in both species, we need to dilute it to 2.5%, and then be used for irrigation, the same trend needs to be followed by Na₂SO₄ concentrations (X_{NS-LT}) and blend of salts (X_{MNC-LT}) and (X_{MNS-LT}), respectively. Therefore, irrigation effects using diluted salted water equivalent to salt stress levels could be predicted through electrophysiological properties.

Table 7. Effect of salt stress and salt stress subsequently re-watering on leaf density of *O. violaceus* and *B. napus*.

Treatments/ re-watering (g L ⁻¹)	<i>L_{TO}. violaceus</i>		<i>L_{TB}. napus</i>	
	Salt stress (cm × 10 ⁻³)	Re-watering (cm × 10 ⁻³)	Salt stress (cm × 10 ⁻³)	Re-watering (cm × 10 ⁻³)
Control	121.0 ^a	131.0 ^a	156.0 ^a	155.0 ^a
NC1/NC1^0	94.0 ^b	106.0 ^b	106.0 ^b	113.0 ^b
NC2/NC2^1	75.6 ^d	101.0 ^c	75.0 ^d	96.6 ^c
NC3/NC3^2	22.6 ^e	36.6 ^d	16.0 ^f	29.4 ^e
NS1/NS1^0	88.3 ^c	102.0 ^c	101.0 ^c	113.0 ^b
NS2/NS2^1	73.9 ^d	101.0 ^c	71.7 ^e	92.9 ^d
NS3/NS3^2	22.7 ^e	37.5 ^d	15.3 ^f	28.8 ^e
BS1/BS1^0	19.7 ^f	29.4 ^f	12.8 ^g	26.9 ^f
BS2/BS2^0	20.2 ^f	31.6 ^e	14.1 ^g	27.4 ^f
BS3/BS3^0	73.3 ^d	102.0 ^c	70.3 ^e	92.1 ^d

Notes: The means ± SE (n = 5) followed by different letter in the columns indicated the leaf density of both species under salt stress followed by re-watering with significant difference at p ≤ .05, according to one-way ANOVA.

NC1: 2.5 g L⁻¹ NaCl, NC2: 5.0 g L⁻¹ NaCl, NC3: 10.0 g L⁻¹ NaCl; NS1: 2.5 g L⁻¹ Na₂SO₄, NS2: 5.0 g L⁻¹ Na₂SO₄, NS3: 10.0 g L⁻¹ Na₂SO₄; BS1: 2.5 g L⁻¹ NaCl + 10.0 g L⁻¹ Na₂SO₄; BS2: 10.0 g L⁻¹ NaCl + 2.5 g L⁻¹ Na₂SO₄; BS3: 5.0 g L⁻¹ NaCl + 5.0 g L⁻¹ Na₂SO₄ and 0 as control. NC1^0: 2.5 g L⁻¹ NaCl → 0 g L⁻¹ NaCl, NC2^1: 5.0 g L⁻¹ NaCl → 2.5 g L⁻¹ NaCl, NC3^2: 10.0 g L⁻¹ NaCl → 5.0 g L⁻¹ NaCl, NS1^0: 2.5 g L⁻¹ Na₂SO₄ → 0 g L⁻¹ Na₂SO₄, NS2^1: 5.0 g L⁻¹ Na₂SO₄ → 2.5 g L⁻¹ Na₂SO₄, NS3^2: 10.0 g L⁻¹ Na₂SO₄ → 5.0 g L⁻¹ Na₂SO₄, BS1^0, BS2^0 and BS3^0 re-watered with control

Restoration of *O. violaceus* and *B. napus* through re-watering

Re-watering had a better effect on *O. violaceus* and *B. napus* and plants successfully recovered. It showed the tolerance and maintenance of plants up to some threshold levels. A substantial increase occurred even during re-watering of *O. violaceus* in PH, SD and LA from low (NC1^0 and NS1^0) to moderate levels (NC2^1, NS2^1 and BS3^0) (Tables 1–3). The increments of PH, SD and LA were found more after re-watering in *O. violaceus* rather than *B. napus*, due to the increase in C_P and recovery of Ψ. It indicates that the water-holding capability of the leaf cell of *O. violaceus* was better to maintain leaf WC. A little contribution was found in this research direction by Flexas et al. (2009), focusing on photosynthetic acclimation followed by re-watering to induce recovery of the plant. In the previous study, *O. violaceus* was found to be shade-tolerant as compared to other Brassicaceae species because its photosynthetic activity was increased by regulation of higher CA activity (Wu et al. 2011). Though, *O. violaceus* exhibited better ability to resist salt stress than *B. napus* after re-watering.

Table 8. Model between salt concentrations and physiological capacitance and leaf tensity.

Species	Model type	Model equation	R ²	n	p
<i>O. violaceus</i>	X _{NC-CP}	CP = 77.03 exp ^{-0.16X_c}	0.9757	12	<.0001
	X _{NC-LT}	LT = 125.46 exp ^{-0.13X_c}	0.9493	12	<.0001
	X _{NS-CP}	CP = 76.15 exp ^{-0.17X_c}	0.9822	12	<.0001
	X _{NS-LT}	LT = 123.85 exp ^{-0.13X_c}	0.9581	12	<.0001
<i>B. napus</i>	X _{NC-CP}	CP = 96.30 exp ^{-0.19X_c}	0.9784	12	<.0001
	X _{NC-LT}	LT = 159.10 exp ^{-0.17X_c}	0.9774	12	<.0001
	X _{NS-CP}	CP = 95.40 exp ^{-0.21X_c}	0.9864	12	<.0001
	X _{NS-LT}	LT = 157.91 exp ^{-0.18X_c}	0.9828	12	<.0001

Notes: X_{NC-LT}: Relationship between NaCl concentrations and leaf tensity; X_{NC-CP}: Relationship between NaCl concentrations and physiological capacitance; X_{NS-LT}: Relationship between Na₂SO₄ concentrations and leaf tensity; X_{NS-CP}: Relationship between Na₂SO₄ concentrations and physiological capacitance.

Salinity affects the leaf characteristic of the plant by inhibiting leaf expansion and elongation (Marcelis & Van Hooijdonk 1999; Qiu et al. 2007). A considerable reduction was observed under high stress levels even during re-watering in growth development of *O. violaceus* and *B. napus* because of no recovery in C_p and unstable status of Ψ (Tables 4 and 6). It also specifies that the ability of the plant to hold water by building leaf cell was not good to maintain water status of leaves. Later T_L of *O. violaceus* and *B. napus* was sustained and recovered at medium stress levels of both single and blend of salts as compared with high stress levels. It is because of restorability of T_L, which can be used to illustrate the water status of the plant and the capacity of holding water. The results in Table 7 show that the maximum increase found under medium stress levels in T_L of *O. violaceus* indicates the maximum recovery of T_L and the ability of *O. violaceus* to hold water. While at the same time recovery of leaf tensity in *B. napus* was also found to be maximum under low-medium stress levels, but the water-holding capability of *B. napus* was not better as was found in *O. violaceus* after re-watering. Though, under moderate stress, T_L directly reflected the ability to resist stress in terms of water loss, whereas salt stress subsequent re-watering reflected the ability to resist stress after dehydration. Thus, this condition exposed the threshold flexibility and more tolerance of *O. violaceus* under moderate levels after re-watering, because of the mechanisms involved in osmotic tolerance related to water availability to sustain leaf cell's energy demands for growth.

Even in the re-watering phase, T_L was still damaged at high concentration levels. This indicates the unhealthy status of plants due to limited water uptake movements, and poor water-holding ability. Plants tolerance to salt stress differs with the degree at which salt reached toxic levels in leaves because of the Ψ difference between the atmosphere and

leaf cells (Carillo et al. 2011). It was hard for both *O. violaceus* and *B. napus* to be recovered from the excessive salt stresses. So, our results agreed with Brodribb and Holbrook (2005), who described that at 50% collapse of cells, it is difficult for a plant to recover after re-watering. Consequently, *O. violaceus* and *B. napus* were thought to be the species with single and blend of salts tolerance flexibility under moderate stress conditions. So, re-watering of saline water should be done under moderate levels. The moderate stress level of both single and blended salts was considered the best for threshold tolerance and production of *O. violaceus* and *B. napus* under saline condition. The effect of salinity in both *O. violaceus* and *B. napus* was reduced. However, in both *O. violaceus* and *B. napus* negligible reduction was observed, which might be attributed to the dilution of saline irrigation (re-watering) and by the combination of salts. Furthermore, the toxic effect of blended salts at BS3^{^0} was found to be the same as it is noted under NC2^{^1} and NS2^{^1} concentration levels of single salts. Hence, *B. napus* can be grown under moderate soil salinity because of its salt tolerance adaptability nature and *O. violaceus* can be grown where saline water resources are existing because of its restoration after re-watering under moderate conditions.

Conclusion

It is concluded that salinity affects the growth of plants by disturbing electrophysiological characteristics of *O. violaceus* and *B. napus*, but the response was different at different salt levels. The maximum recovery of C_p and T_L at medium stress levels after re-watering shows better water-holding capability and growth restorability of *O. violaceus* than *B. napus*. Because at 10% salt concentration, salt stress was too severe to be re-watered, and the irrigation effects using diluted salted water (2.5%) at 5% salt stress level in *O. violaceus* and *B. napus* were equivalent to the effects using 1.66% salted water and 2.88% salted water, respectively, which were equal to the salt stress level between 0% and 2.5%. Accordingly, when the concentration of salted water was 2.5% for both species, it could be used for irrigation directly, but when the concentration of salted water was 5% in both species, we need to dilute it to 2.5%, and then use it for irrigation, the same trend needs to be followed for Na₂SO₄ concentrations (X_{NS-LT}) and blend of salts (X_{MNC-LT}) and (X_{MNS-LT}), respectively. However, from the results (Table 9), it is clear that the dilution point exists in NC2, NS2 and BS3, 5–2.5% for single NaCl and Na₂SO₄ and 10–0.0% for blend of salts. Hence, prediction of dilution point of saline irrigation based on electrophysiological properties could be helpful to maintain crop productivity,

Table 9. Equivalent salt stress levels of both species based on physiological capacitance and leaf tensity.

Species	Concentration (%) single salts/mixed salts	X _{NC-LT} (%)	X _{NC-CP} (%)	X _{MNC-LT} (%)	X _{MNC-CP} (%)	X _{NS-LT} (%)	X _{NS-CP} (%)	X _{MNS-LT} (%)	X _{MNS-CP} (%)
<i>O. violaceus</i>	0–0/0–0	–0.33	–0.44	–0.33	–0.44	–0.42	–0.48	–0.42	–0.48
	2.5–0/12.5–0	1.30	1.20	11.15	13.40	1.45	1.28	10.76	12.61
	5–2.5/12.5–0	1.66	1.74	10.60	12.17	1.53	1.78	10.22	11.46
	10–5/10–0	9.47	9.75	1.60	1.94	8.94	9.21	1.45	1.77
<i>B. napus</i>	0–0/0–0	0.15	–0.09	0.15	–0.09	0.10	–0.12	0.10	–0.12
	2.5–0/12.5–0	1.97	1.67	10.26	13.39	1.84	1.69	9.75	12.11
	5–2.5/12.5–0	2.88	2.98	10.16	12.01	2.92	3.01	9.65	10.86
	10–5/10–0	9.75	10.65	3.15	2.95	9.38	10.57	2.97	2.81

Notes: X_{NC-LT}: Relationship between NaCl concentrations and leaf tensity; X_{NC-CP}: Relationship between NaCl concentrations and physiological capacitance; X_{MNC-LT}: Relationship between NaCl concentrations in blend of salts and leaf tensity; X_{MNC-CP}: Relationship between NaCl concentrations in blend of salts and physiological capacitance; X_{NS-LT}: Relationship between Na₂SO₄ concentrations and leaf tensity; X_{NS-CP}: Relationship between Na₂SO₄ concentrations and physiological capacitance; X_{MNS-LT}: Relationship between Na₂SO₄ concentrations in blend of salts and leaf tensity; X_{MNS-CP}: Relationship between Na₂SO₄ concentrations in blend of salts and physiological capacitance.

reduce irrigation cost, save water resources and could be a better approach to utilize existing water resources.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This study were supported by the National Natural Science Foundation of China [grant number 31301243]; the National Key Basic Research Program (973) special projects [2013CB956701]; the Priority Academic Program Development (PAPD) of Jiangsu Higher Education Institutions, the research foundation for introduce talents of Jiangsu University[13JDG030].

References

- Abdul Qados AMS. 2011. Effect of salt stress on plant growth and metabolism of bean plant *Vicia faba* (L.). JSSAS. 10:7–15.
- Álvarez S, Gómez-Bellot MJ, Castillo M, Bañón S, Sánchez-Blanco MJ. 2012. Osmotic and saline effect on growth, water relations, and ion uptake and translocation in *Phlomis purpurea* plants. Environ Exp Bot. 78:138–145.
- Aref F. 2013. Effect of saline irrigation water on yield and yield components of rice (*Oryza sativa* L.). Afr J Biotechnol. 12:3503–3513.
- Ashraf M, McNeilly T. 2004. Salinity tolerance in brassica oilseeds. Crit Rev Plant Sci. 23:157–174.
- Benincasa P, Pace R, Quinet M, Lutts S. 2013. Effect of salinity and priming on seedling growth in rapeseed (*Brassica napus* var *oleifera* Del.). Acta Sci Agron. 35:479–486.
- Brodribb TJ, Holbrook NM. 2005. Water stress deforms tracheids peripheral to the leaf vein of a tropical conifer. Plant Physiol. 137:1139–1146.
- Carillo P, Annunziata MG, Pontecorvo G, Fuggi A, Woodrow P. 2011. Salinity stress and salt tolerance. In: A. Shanker, editor. Abiotic stress plants-mech adapt. Rijeka, Croatia: In Tech; p. 21–38.
- de Oliveira AB, Gomes-Filho E, Alencar NLM, Akinci S. 2013. Comparison between the water and salt stress effects on plant growth and development. In: A. Sener, editor. Response of organisms to water stress. Rijeka, Croatia: InTech; p. 67–94.
- Flexas J, Barón M, Bota J, Ducruet J-M, Gallé A, Galmés J, Jiménez M, Pou A, Ribas-Carbó M, Sajnani C, et al. 2009. Photosynthesis limitations during water stress acclimation and recovery in the drought-adapted *Vitis* hybrid Richter-110 (*V. berlandieri* × *V. rupestris*). J Exp Bot. 60:2361–2377.
- Fu A, Chen Y, Li W. 2010. Analysis on the change of water potential of *Populus euphratica* Oliv. and *P. Russkii* Jabl under different irrigation volumes in temperate desert zone. Chin Sci Bull. 55:965–972.
- García-Sánchez F, Jifon JL, Carvajal M, Syvertsen JP. 2002. Gas exchange, chlorophyll and nutrient contents in relation to Na⁺ and Cl⁻ accumulation in ‘sunburst’ mandarin grafted on different rootstocks. Plant Sci. 162:705–712.
- Gul H, Ahmad R. 2004. Effect of different irrigation intervals on growth of canola (*Brassica napus* L.) under different salinity levels. Pak J Bot. 36:359–372.
- Hoagland DR, Arnon DI. 1950. The water-culture method for growing plants without soil. Circ Calif Agric Exp Sta. 347:1–32.
- Irigoyen J, Einerich D, Sánchez-Díaz M. 1992. Water stress induced changes in concentrations of proline and total soluble sugars in nodulated alfalfa (*Medicago sativa*) plants. Physiol Plant. 84:55–60.
- Jamil M, Bashir S, Anwar S, Bibi S, Bangash A, Ullah F, Rha ES. 2012. Effect of salinity on physiological and biochemical characteristics of different varieties of rice. Pak J Bot. 44:7–13.
- Koide RT, Robichaux RH, Morse SR, Smith CM. 1989. Plant water status, hydraulic resistance and capacitance. In: Plant physiological ecology: field methods and instrumentation. Dordrecht: Springer; p. 161–183.
- Mansour MMF, Salama KHA. 2004. Cellular basis of salinity tolerance in plants. Environ Exp Bot. 52:113–122.
- Marcelis L, Van Hooijdonk J. 1999. Effect of salinity on growth, water use and nutrient use in radish (*Raphanus sativus* L.). Plant Soil. 215:57–64.
- Masidur Alam M, Hayat S, Ali B, Ahmad A. 2007. Effect of 28-homobrassinolide treatment on nickel toxicity in *Brassica juncea*. Photosynthetica. 45:139–142.
- Meinzer F, James S, Goldstein G, Woodruff D. 2003. Whole-tree water transport scales with sapwood capacitance in tropical forest canopy trees. Plant Cell Environ. 26:1147–1155.
- Munns R. 2002. Comparative physiology of salt and water stress. Plant Cell Environ. 25:239–250.
- Munns R, Tester M. 2008. Mechanisms of salinity tolerance. Annu Rev Plant Biol. 59:651–681.
- Niu G, Rodriguez DS, Aguiniga L. 2008. Effect of saline water irrigation on growth and physiological responses of three rose rootstocks. HortScience. 43:1479–1483.
- Purty RS, Kumar G, Singla-Pareek SL, Pareek A. 2008. Towards salinity tolerance in Brassica: an overview. Physiol Mol Biol Plants. 14:39–49.
- Qiu D, Lin P, Guo S. 2007. Effects of salinity on leaf characteristics and CO₂/H₂O exchange of *Kandelia candel* (L.) Druce seedlings. J For Sci. 53:13–19.
- Roy SJ, Negrão S, Tester M. 2014. Salt resistant crop plants. Curr Opin Biotech. 26:115–124.
- SEMÍZ GD, Suarez DL. 2015. Tomato salt tolerance: impact of grafting and water composition on yield and ion relations. Turk J Agric For. 39:876–886.
- Siringam K, Juntawong N, Cha-um S, Kirdmanee C. 2011. Salt stress induced ion accumulation, ion homeostasis, membrane injury and sugar contents in salt-sensitive rice (*Oryza sativa* L. spp. indica) roots under isoosmotic conditions. Afr J Biotechnol. 10:1340–1346.
- Su J, Wu S, Xu Z, Qiu S, Luo T, Yang Y, Chen Q, Xia Y, Zou S, Huang B-L. 2013. Comparison of salt tolerance in brassicas and some related species. AJPS. 4:1911–1917.
- Tester M, Bacic A. 2005. Abiotic stress tolerance in grasses. From model plants to crop plants. Plant Physiol. 137:791–793.
- Vogt UK, Losch R. 1999. Stem water potential and leaf conductance: A comparison of *Sorbus aucuparia* and *Sambucus nigra*. Phys Chem Earth Pt B: Hydrology, Oceans and Atmosphere. 24:121–123.
- Wang X, Geng S, Ri Y-J, Cao D, Liu J, Shi D, Yang C. 2011. Physiological responses and adaptive strategies of tomato plants to salt and alkali stresses. Sci Hortic-Amsterdam. 130:248–255.
- Wu Y, Shi Q, Wang K, Li P, Xing D, Zhu Y, Song Y. 2011. An electrochemical approach coupled with Sb microelectrode to determine the activities of carbonic anhydrase in the plant leaves. In: D. Zheng, editor. Future intelligent information systems. Berlin: Springer-Verlag; p. 87–94.
- Yao X, Yang R, Zhao F, Wang S, Li C, Zhao W. 2016. An analysis of physiological index of differences in drought tolerance of tomato rootstock seedlings. J Plant Biol. 59:311–321.
- Zapryanova N, Atanassova B. 2009. Effects of salt stress on growth and flowering of ornamental annual species. Biotechnol Biotech Equip. 23:177–179.
- Zhang M, Wu Y, Xing D, Zhao K, Yu R. 2015. Rapid measurement of drought resistance in plants based on electrophysiological properties. ASABE. 58:1441–1446.
- Zhang X, Wang Z, Zhang X, Li M, Zou J. 2012. Effects of heavy metals and saline-alkali on growth, physiology and biochemistry of *Orychophragmus violaceus*. Agric Sci Technol (农业科学与技术). 13:1478–1483.
- Zollinger N, Koenig R, Cerny-Koenig T, Kjelgren R. 2007. Relative salinity tolerance of intermountain western United States native herbaceous perennials. HortScience. 42:529–534.